

The H-Boltzmann's Theorem for Magnetic Systems

S. Malinowski

University of Łódź
Department of Theoretical Physics
ul. Pomorska 149/153
90-236 Łódź, Poland

June 1995

Abstract

In a time-reversal non-invariant system the entropy production, dS/dt , may be negative. Close to the critical point of medium it is positive: larger or smaller as compared with one above critical point depending on whether the transitions from lower states to higher states of the system are more (less) probable than the reverse transitions. PACS numbers: 05.60.+w, 05.70.Fh, 05.70.Ln

The entropy S is defined [1] as follows:

$$S = -k \sum_{n=1}^{\infty} P_n \ln P_n \quad (1)$$

where k is Boltzmann's constant and

$$P_n = \frac{\exp(-E_n/kT)}{\sum_{m=1}^{\infty} \exp(-E_m/kT)} \quad (2)$$

is the Gibbs distribution of the energy levels E_n . To clarify the considerations we will use the following arrangement for the energy levels:

$$E_1 \leq E_2 \leq \dots \leq E_{\infty}. \quad (3)$$

Hence, for the probability distributions we obtain the relations:

$$1 \geq P_1 \geq P_2 \geq \dots \geq P_{\infty} \geq 0. \quad (4)$$

The time derivative of entropy is equal to

$$\frac{dS}{dt} = -k \sum_{n=1}^{\infty} \frac{dP_n}{dt} \ln P_n \quad (5)$$

because

$$\frac{d}{dt} \sum_{n=1}^{\infty} P_n = 0.$$

From Boltzmann's equation we have

$$\frac{dP_n}{dt} = \sum_{m=1}^{\infty} (W_{m,n}P_m - W_{n,m}P_n), \quad (6)$$

where $W_{m,n}$ is the transition probability from the state ψ_m (level E_m) to the state ψ_n (level E_n). Hence, for dS/dt we have the following equivalent formulae:

$$\frac{dS}{dt} = -k \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} (W_{m,n}P_m \ln P_n - W_{n,m}P_n \ln P_n) \quad (7a)$$

$$= k \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} W_{n,m}P_n \ln \frac{P_n}{P_m} \quad (7b)$$

$$= k \sum_{n=1}^{\infty} \sum_{i=n+1}^{\infty} (W_{n,n+i} P_n - W_{n+i,n} P_{n+i}) \ln \frac{P_n}{P_{n+i}} \quad (7c)$$

Let us compare the entropy production in the same sample before and after breaking its time-inversion symmetry. Consider the electronic system in the temperature T_1 above, $T_c < T_1$, and T_2 below, $T_2 < T_c$, a critical temperature T_c is an example of such medium.

In temperature T_1 the matrix of transition probabilities is a symmetric one:

$$W_{n,n+i}(0) = W_{n+i,n}(0) \equiv W_{n,n+i}^{(s)}, \quad (8)$$

while in temperature T_2 it is not symmetric and moreover it is some function of spontaneous magnetization $\mathbf{M}$ of the sample:

$$W_{n,n+i}(\mathbf{M}) = W_{n,n+i}^{(s)}(\mathbf{M}) + W_{n,n+i}^{(a)}(\mathbf{M}). \quad (9)$$

Hence, we have

$$W_{n+i,n}(\mathbf{M}) = W_{n,n+i}^{(s)}(\mathbf{M}) - W_{n,n+i}^{(a)}(\mathbf{M}). \quad (10)$$

The time-inversion operator transforms $W_{n,n+i}(\mathbf{M})$ into $W_{n+i,n}(-\mathbf{M})$. It means that the $W^{(s)}$ is an even function of $\mathbf{M}$,

$$W_{n,n+i}^{(s)}(\mathbf{M}) = W_{n,n+i}^{(s)}(-\mathbf{M}) \quad (11)$$

and $W^{(a)}$ - an odd one,

$$W_{n,n+i}^{(a)}(\mathbf{M}) = -W_{n,n+i}^{(a)}(-\mathbf{M}). \quad (12)$$

The other properties are following:

$$\lim_{\mathbf{M} \rightarrow 0} W_{n,n+i}^{(s)}(\mathbf{M}) \rightarrow W_{n,n+i}^{(s)}(0), \quad (13)$$

$$\lim_{\mathbf{M} \rightarrow 0} W_{n,n+i}^{(a)}(\mathbf{M}) = 0, \quad (14)$$

$$|W_{n,n+i}^{(a)}(\mathbf{M})| \leq W_{n,n+i}^{(s)}(\mathbf{M}). \quad (15)$$

The latter follows from the definition of the $W_{n,n+i}$.

The $W_{n,n+i}^{(a)}(\mathbf{M})$ may be positive or negative. If it is positive:

$$W_{n,n+i}^{(a)}(\mathbf{M}) > 0 \quad (16)$$

then the probability of transition from the lower state ψ_n to the higher state ψ_{n+i} is larger than the probability of the reverse transition. If it is negative:

$$W_{n,n+i}^{(a)}(\mathbf{M}) < 0 \quad (17)$$

then we are dealing with the reverse relation between the probabilities of transition.

Inserting the expressions (9) and (10) to the formula (7c) we obtain

$$\begin{aligned} \frac{dS(\mathbf{M})}{dt} &= k \sum_{n=1}^{\infty} \sum_{i=n+1}^{\infty} [W_{n,n+1}^{(s)}(\mathbf{M})(P_n - P_{n+i}) + W_{n,n+i}^{(a)}(\mathbf{M})(P_n + P_{n+i})] \ln \frac{P_n}{P_{n+i}} \\ &= \frac{dS(\mathbf{M})_{\text{inv}}}{dt} + \frac{dS(\mathbf{M})_{\text{noninv}}}{dt} \end{aligned} \quad (18)$$

where the invariant part of $dS(\mathbf{M})/dt$ with respect to the time inversion is always positive:

$$\frac{dS(\mathbf{M})_{\text{inv}}}{dt} = k \sum_{n=1}^{\infty} \sum_{i=n+1}^{\infty} W_{n,n+i}^{(s)}(\mathbf{M})(P_n - P_{n+i}) \ln \frac{P_n}{P_{n+i}} > 0, \quad (19)$$

and the noninvariant part

$$\frac{dS(\mathbf{M})_{\text{noninv}}}{dt} = k \sum_{n=1}^{\infty} \sum_{i=n+1}^{\infty} W_{n,n+i}^{(a)}(\mathbf{M})(P_n + P_{n+i}) \ln \frac{P_n}{P_{n+i}}$$

may acquire, by ineq. (16) and (17) an arbitrary sign.

IN CONCLUSION: Far apart from the critical point of a sample, when $W^{(a)}$ may achieve the absolute values comparable with the values of $W^{(s)}$ and almost all $W_{n,n+i}^{(a)}(\mathbf{M})$ are negative the entropy production may be negative

$$\frac{dS(\mathbf{M})}{dt} < 0. \quad (20)$$

Close to the critical point, due to the properties (13) and (14), the entropy production is positive and for some samples is larger than the one above critical point,

$$\frac{dS(\mathbf{M})}{dt} > \frac{dS(0)}{dt} > 0 \quad (21)$$

and for the others is smaller

$$\frac{dS(0)}{dt} > \frac{dS(\mathbf{M})}{dt} > 0. \quad (22)$$

Transport phenomena are examples of irreversible processes. The formulae (21) and (22) translated to the behaviour of the electric resistivity ρ in the neighbourhood of critical point lead to the inequality

$$\rho(\mathbf{M}) < \rho(0) \quad (23)$$

and

$$\rho(0) > \rho(\mathbf{M}), \quad (24)$$

respectively, which is well-known to experimentators [2] (see the papers listed in [2]). In this situation the above generalization of Boltzmann's theorem can be treated as the additional interpretation of phase transitions. The explicit form of the transition probability $W_{n,m}$ satisfying the requirement (9) will be subject of the next paper.

Acknowledgement

I wish to thank Professor P. Kosiski for fruitful discussions and reading the manuscript

References

- [1] R.P. Feynman, Statistical Mechanics. A Set of Lectures, W.A. Benjamin, 1972.
- [2] S. Malinowski, Acta Phys. Pol. **A77**, 569 (1990).